

1. Let $f(x) = x^3 + 9x$ on $[1, 3]$.

<p>(a) What is the average rate of change in f on $[1, 3]$?</p> $\begin{aligned}\text{average rate} &= \frac{f(3) - f(1)}{3 - 1} \\ &= \frac{54 - 10}{2} \\ &= 22\end{aligned}$	<p>+1 Use difference quotient +1 Answer is 22</p>
<p>(b) On what interval(s) is $f'(x) > 0$?</p> $\begin{aligned}f'(x) &= 3x^2 + 9 \\ &= 3(x^2 + 3)\end{aligned}$ <p>Since $x^2 + 3 > 0$ for all x, $f'(x) > 0$ for all x. Thus $f'(x) > 0$ on the interval $[1, 3]$.</p>	<p>+1 $f'(x) = 3x^2 + 9$ +1 Justify why $f'(x) > 0$ for all x. +1 The interval $[1, 3]$</p>
<p>(c) Find a value c in the interval $[1, 3]$ such that the average rate of change in f over the interval $[1, 3]$ equals $f'(c)$.</p> $\begin{aligned}f'(c) &= 3c^2 + 9 \\ 22 &= 3c^2 + 9 \\ 13 &= 3c^2 \\ c^2 &= \frac{13}{3} \\ c &= \sqrt{\frac{13}{3}} \\ &\approx 2.082\end{aligned}$	<p>+1 $f'(c) = 3c^2 + 9$ +1 Set $f'(c)$ equal to average rate of change from (a) or equal to 22 +2 $c = \sqrt{\frac{13}{3}} \approx 2.082$ (-1 if $-\sqrt{\frac{13}{3}}$)</p>


2. Let $f(x) = \sin x - \cos x$ on $[0, \pi]$.

<p>(a) What is the average rate of change in f on $[0, \pi]$?</p> $\begin{aligned} \text{average rate} &= \frac{f(\pi) - f(0)}{\pi - 0} \\ &= \frac{\sin(\pi) - \cos(\pi) - (\sin(0) - \cos(0))}{\pi} \\ &= \frac{0 - (-1) - (0 - 1)}{\pi} \\ &= \frac{2}{\pi} \end{aligned}$	<p>+1 Use difference quotient +1 Answer is $\frac{2}{\pi}$</p>
<p>(b) On what interval(s) is $f'(x) > 0$?</p> <p>$g'(x) = \cos x + \sin x$ on $[0, \pi]$.</p> <p>Since the coordinate for angle x on the unit circle is $(\cos x, \sin x)$, $g'(x) = \cos x + \sin x$ is the sum of the coordinate values for angle x.</p> <p>In the first quadrant, both coordinate values are positive so $g'(x) > 0$ for $0 < x < \frac{\pi}{2}$. At $x = 0$ and $x = \frac{\pi}{2}$, the coordinates are $(1, 0)$ and $(0, 1)$, respectively, so $g'(x) > 0$ at those values also.</p> <p>In the second quadrant, $\cos x < 0$. Additionally, $\sin x > \cos x$ for $\frac{\pi}{2} < x < \frac{3\pi}{4}$. Therefore, $\cos x + \sin x > 0$ on $\frac{\pi}{2} < x < \frac{3\pi}{4}$.</p> <p>Thus $g'(x) > 0$ for $0 < x < \frac{3\pi}{4}$.</p>	<p>+1 $g'(x) = \cos x + \sin x$ +1 $g'(x) > 0$ for $0 \leq x \leq \frac{\pi}{2}$ with valid argument +1 $g'(x) > 0$ for $\frac{\pi}{2} < x < \frac{3\pi}{4}$ with valid argument</p>
<p>(c) Find a value c in the interval $[0, \pi]$ such that the average rate of change in f over the interval $[0, \pi]$ equals $f'(c)$.</p> $\cos c + \sin c = \frac{2}{\pi}$ <p>Solving with technology, $x \approx 1.889$.</p>	<p>+1 $f'(c) = \cos c + \sin c$ +1 Set $f'(c)$ equal to average rate of change from (a) or equal to 22 +2 $x \approx 1.889$</p>

4. Let $f(x) = \frac{2x}{x^2 + 1}$ on $[-5, 5]$.

<p>(a) Find the critical value(s) of f.</p> $f'(x) = \frac{2(x^2 + 1) - 2x(2x)}{(x^2 + 1)^2}$ $= \frac{-2x^2 + 2}{(x^2 + 1)^2}$ $0 = \frac{-2(x-1)(x+1)}{(x^2 + 1)^2}$ $x = \pm 1$	<p>+1 $f'(x) = \frac{-2x^2 + 2}{(x^2 + 1)^2}$</p> <p>+1 Set $f'(x) = 0$</p> <p>+1 $x = \pm 1$</p>
<p>(b) What are the coordinates of the relative extrema? Justify your conclusion.</p> <p>We evaluate the derivative on both sides of the critical values and draw the following conclusions.</p> $f'(x) < 0 \text{ for } x < -1$ $f'(-1) = 0$ $f'(x) > 0 \text{ for } 0 < x < 1$ $f'(1) = 0$ $f'(x) < 0 \text{ for } x > 1$ <p>By the First Derivative Test, a relative minimum occurs at $x = -1$ and a relative maximum occurs at $x = 1$. The corresponding coordinates are $(-1, -1)$ and $(1, 1)$, respectively.</p>	<p>+1 Show f' changes from negative to positive at $x = -1$.</p> <p>+1 Show f' changes from positive to negative at $x = 1$.</p> <p>+1 A relative minimum is $(-1, -1)$</p> <p>+1 A relative maxima is $(1, 1)$.</p>
<p>(c) What is $\lim_{x \rightarrow \infty} f(x)$? Justify your solution.</p> <p>As f approaches infinity, $f(x) = \frac{2x}{x^2 + 1}$ may be closely approximated by $y = \frac{2x}{x^2} = \frac{2}{x}$. We know that $\lim_{x \rightarrow \infty} \frac{2}{x} = 0$. Therefore,</p> $\lim_{x \rightarrow \infty} f(x) = 0.$	<p>+1 $\lim_{x \rightarrow \infty} f(x) = 0$</p> <p>+1 Justification</p>


6. The graph of $f(t)$ is shown in the figure below.



//Comp: Equation is $s(t) = (t-1)(t-3)(t+2)^2$


<p>(a) Estimate from the graph the interval(s) on which f' is positive. Explain how you know.</p> <p>Where the graph is increasing, f' is positive. The graph appears to be increasing on the intervals $-2 < t < -0.5$ and $2.5 < t < 4$. Thus f' is positive on those intervals.</p>	<p>+1 $-2 < t < -0.5$ +1 $2.5 < t < 4$ +1 Where the graph is increasing, f' is positive.</p>
<p>(b) Estimate from the graph the interval(s) on which f'' is positive. Explain how you know.</p> <p>Where the graph is concave up, f'' is positive. The graph appears to be concave up on the intervals $-4 < t < -1$ and $1 < t < 4$. Thus f'' is positive on those intervals.</p>	<p>+1 $-4 < t < -1$ +1 $1 < t < 4$ +1 Where the graph is concave up, f'' is positive.</p>
<p>(c) Estimate from the graph the values of t where $f'(t) = 0$. Explain how you know.</p> <p>Since the graph of f is continuous and differentiable, we need only determine where the graph has horizontal tangent lines. This will occur at a relative extrema. The relative extrema appear to occur at $t = -2, -0.5$, and 2.5.</p>	<p>+1 $t = -2, -0.5$, and 2.5 +2 Explanation</p>

8. An open rectangular box with square base has a volume of 256 cubic inches.



<p>(a) Determine the equation for the volume and surface area of the box.</p> $V = x^2h \text{ and } A = x^2 + 4xh$	<p>+1 $V = x^2h$ +1 $A = x^2 + 4xh$</p>
<p>(b) What box dimensions minimize surface area?</p> $256 = x^2h$ $h = \frac{256}{x^2}$ $A = 4xh + x^2$ $= 4x\left(\frac{256}{x^2}\right) + x^2$ $= 1024x^{-1} + x^2$ $A' = -1024x^{-2} + 2x$ $0 = -\frac{1024}{x^2} + 2x$ $\frac{1024}{x^2} = 2x$ $1024 = 2x^3$ $x^3 = 512 \Rightarrow x = 8$ <p>The minimum dimensions are 8 inches by 8 inches by 4 inches.</p>	<p>+1 $h = \frac{256}{x^2}$ +1 $A' = -1024x^{-2} + 2x$ +1 $x = 8$ inches +1 $h = 4$ inches +1 verify minimum</p>
<p>(c) With the added restriction that the box height cannot exceed 3 inches, what is the minimum surface area?</p> <p>We know that A is decreasing on the interval $0 < x < 8$ and increasing for $x > 8$. Our objective is to pick x as close to 8 as possible. Since $256 = x^2h$, $x = \sqrt{\frac{256}{h}}$. As h increases, x decreases. Since the maximum value of h is 3, the minimum value of x is $x = \sqrt{\frac{256}{3}} \approx 9.238$. The minimum surface area is $A = \left(\sqrt{\frac{256}{3}}\right)^2 + 4\left(\sqrt{\frac{256}{3}}\right)(3) \approx 196.185$ sq in.</p>	<p>+1 $x = \sqrt{\frac{256}{3}} \approx 9.238$ +1 $A \approx 196.185$ sq in</p>

10. A new pen is to be built between a barn and an existing pen as shown in the figure. The sides of the new pen bordered by the barn and the existing pen will not require any new fence.



<p>(a) Write an equation that represents the amount of fence needed to enclose the new pen.</p> $P = x + 40 + y + 30 + x + y$ $= 2x + 2y + 70$	<p>+1 $P = 2x + 2y + 70$</p>
<p>(b) Write an equation that represents the area of the new pen.</p> $A = (40 + y)(30 + x)$	<p>+1 $A = (40 + y)(30 + x)$</p>
<p>(c) If the new pen needs contain 1444 square feet of area, what pen size minimizes the amount of fence needed?</p> $1444 = (40 + y)(30 + x)$ $x = \frac{1444}{40 + y} - 30$ $x = \frac{1444}{40 + (-2)} - 30$ $= 38 - 30$ $= 8$ $30 + 8 = 38$ $40 + (-2) = 38$ <p>The pen should be 38 feet by 38 feet.</p> $P = 2x + 2y + 70$ $= 2\left(\frac{1444}{40 + y} - 30\right) + 2y + 70$ $= \frac{2888}{40 + y} - 60 + 2y + 70$ $= \frac{2888}{40 + y} + 2y + 10$ $P' = -2888(40 + y)^{-2} + 2$ $0 = -2888(40 + y)^{-2} + 2$ $2 = 2888(40 + y)^{-2}$ $2(40 + y)^2 = 2888$ $(40 + y)^2 = 1444$ $40 + y = 38$ $y = -2$	<p>+1 $x = \frac{1444}{40 + y} - 30$</p> <p>+1 $P = \frac{2888}{40 + y} + 2y + 10$</p> <p>+1 $P' = -2888(40 + y)^{-2} + 2$</p> <p>+1 Set $P' = 0$</p> <p>+1 $y = -2$</p> <p>+1 $x = 8$</p> <p>+1 38 feet by 38 feet</p>